

L^2 -spectral invariants and quasi-crystal graphs

GÁBOR ELEK *

Abstract. Introducing and studying the pattern frequency algebra, we prove the analogue of Lück's approximation theorems on L^2 -spectral invariants in the case of aperiodic order. These results imply a uniform convergence theorem for the integrated density of states as well as the positivity of the logarithmic determinant of certain discrete Schrodinger operators.

AMS Subject Classifications: 81Q10, 46L51

Keywords: quasi-crystal graphs, L^2 -invariants, von Neumann algebras, integrated density of states

*The Alfred Renyi Mathematical Institute of the Hungarian Academy of Sciences, P.O. Box 127, H-1364 Budapest, Hungary. email:elek@renyi.hu, Supported by OTKA Grants T 049841 and T 037846

1 Introduction

The goal of the Oberwolfach-Mini-Workshop “ L^2 -spectral invariants and the integrated density of states” was to unify the point of views and approaches in certain areas of geometry and mathematical physics. The aim of our paper is to make the connection between those fields even more explicit. Let us start with a very brief introduction to L^2 -spectral invariants.

1.1 L^2 -spectral invariants

Let Γ be a countable group and $L^2(\Gamma)$ be the Hilbert-space of the formal sums $\sum_{\gamma \in \Gamma} a_\gamma \cdot \gamma$, where $a_\gamma \in \mathbb{C}$ and $\sum_{\gamma \in \Gamma} |a_\gamma|^2 < \infty$. Notice that Γ unitarily acts on $L^2(\Gamma)$ by $L_\delta(\sum_{\gamma \in \Gamma} a_\gamma \cdot \gamma) = \sum_{\gamma \in \Gamma} a_{\gamma\delta^{-1}} \cdot \gamma$. Hence one can represent the complex group algebra $\mathbb{C}\Gamma$ as bounded operators by left convolutions. The weak closure of $\mathbb{C}\Gamma$ in $B(L^2(\Gamma))$ is the group von Neumann algebra $\mathcal{N}\Gamma$. The group von Neumann algebra has a natural trace:

$$\mathrm{Tr}_\Gamma(A) = \langle A(1), 1 \rangle ,$$

where $1 \in L^2(\Gamma)$ is identified with the unit element of the group. Let $B \in \mathcal{N}\Gamma$ be a positive, self-adjoint element, then by the spectral theorem of von Neumann

$$B = \int_0^\infty \lambda dE_\lambda^B ,$$

where $E_\lambda^B = \chi_{[-\infty, \lambda]}(B)$. We can associate a *spectral measure* μ_B to our operator B by

$$\mu_B[0, \lambda] = \sigma_B(\lambda) = \mathrm{Tr}_\Gamma E_\lambda^B .$$

Note that the jumps of σ_B are associated to the eigenspaces of B . We denote by S_λ^B the orthogonal projection onto the subspace of λ -eigenvectors. One can also consider the Fuglede-Kadison determinant as

$$\det_\Gamma(B) := \exp\left(\int_0^\infty \log \lambda d\mu_B\right) .$$

Now let Γ be a finitely generated amenable group. That is Γ has a Følner-exhaustion of finite subsets $1 \in F_1 \subset F_2 \subset \dots, \cup_{n=1}^\infty F_n = \Gamma$ such that

$$\lim_{n \rightarrow \infty} \frac{|KF_n|}{|F_n|} = 1 ,$$

for any non-empty finite subset $K \subset \Gamma$. Then the following approximation theorem holds:

Statement 1 [1] *For any positive, self-adjoint $B \in \mathbb{C}(\Gamma)$ and $\lambda \geq 0$,*

$$\text{Tr} S_\lambda^B = \lim_{n \rightarrow \infty} \frac{\dim_{\mathbb{C}} \text{Ker}(B_n - \lambda)}{|F_n|},$$

where $B_n = p_{F_n} B i_{F_n}$ and $p_{F_n} : L^2(\Gamma) \rightarrow L^2(F_n)$ is the natural projection operator, $i_{F_n} : L^2(F_n) \rightarrow L^2(\Gamma)$ is the adjoint of p_{F_n} , the natural imbedding operator.

As we shall see in Section 2 the fact that the spectral distribution functions N_{B_n} converge to σ_B uniformly immediately follows from Statement 1. Note that

$$N_{B_n}(\lambda) = \frac{s(\lambda, B_n)}{|F_n|},$$

where $s(\lambda, B_n)$ is the number of eigenvalues of B_n (counted with multiplicities) not greater than λ .

Similar approximation theorem holds for residually finite groups. Let Γ be a finitely generated residually finite group with finite index normal subgroups

$$\Gamma \triangleright N_1 \triangleright N_2 \triangleright \dots, \cap_{k=1}^{\infty} N_k = \{1\}.$$

Statement 2 [5] *Let $B \in \mathbb{Q}\Gamma$ be a positive, self-adjoint element and let $\pi_k(B) = B_k \in \mathbb{Q}(\Gamma/N_k)$ be the associated finite dimensional linear operators, where $\pi_k : \Gamma \rightarrow \Gamma/N_k$ are the quotient maps. Then the distribution function σ_B is the uniform limit of the spectral distribution functions N_{B_k} .*

Note that the rationality assumption is crucial in the proof of Statement 2. For the Fuglede-Kadison determinant one has the following result:

Statement 3 [5] *If Γ is amenable or residually finite, and $B \in \mathbb{Q}\Gamma$ as above then $\det_{\Gamma}(B) \geq 1$, that is $\int_0^{\infty} \log \lambda d\mu_B \geq 0$.*

1.2 Quasi-crystal graphs

Let $G(V, E)$ be an infinite, connected graph with bounded vertex degrees, which is amenable in the sense that there exists a Følner-sequence $\{F_n\}_{n=1}^\infty$ of finite subsets in V such that

$$\lim_{n \rightarrow \infty} \frac{|\partial F_n|}{|F_n|} = 1,$$

where

$$\partial F_n := \{p \in F_n \mid \text{there exists } q \notin F_n, (p, q) \in E\}.$$

We shall consider those graphs that exhibit a certain aperiodic order. First we need some definitions.

The r -pattern of a vertex $x \in V$ is the graph automorphism class of the rooted ball around x . That is $x, y \in V$ have the same r -pattern if there exists a graph isomorphism ϕ between the balls $B_r(x)$ and $B_r(y)$ such that $\phi(x) = y$. Denote by $P_r(G)$ the finite set of all possible r -patterns in G . We say that G is an *abstract quasicrystal graph* (AQ-graph) if for any $\alpha \in P_r(G)$ there exists a *frequency* $P(\alpha)$ such that

- For *any* Følner-sequence $\{Q_n\}_{n=1}^\infty$:

$$\lim_{n \rightarrow \infty} \frac{|Q_n^\alpha|}{|Q_n|} = P(\alpha),$$

where $Q_n^\alpha \subseteq Q_n$ is the set of vertices in Q_n having the r -pattern α .

Of course, the Cayley-graphs of finitely generated groups are AQ-graphs. Also, the dual graph of the Penrose tilings and in general, graphs defined by nice Delone sets considered in [3], [4] are AQ-graphs without translational symmetries. We can construct random AQ-graphs as well. Let G be the standard lattice graph in the two-dimensional plane. Consider the Bernoulli space $\Omega = \prod_G \{0, 1\}$. For each $\omega \in \Omega$, G_ω is the following graph. If $\omega(a, b) = 0$ then we add a new edge $((a, b), (a + 1, b + 1))$ to the lattice. Then for almost all $\omega \in \Omega$, G_ω is an AQ-graph.

The algebra that plays the role of the group algebra is the *algebra of pattern invariant matrices* P_G . A pattern-invariant matrix is a function $A : V \times V \rightarrow \mathbb{C}$ satisfying the following properties:

- $A(x, y) = 0$ if $d_G(x, y) > r_A$, where d_G is the shortest path distance on the vertices. The value r_A , the propagation of A , depends only on A .
- $A(x, y) = A(\phi(x), \phi(y))$ if ϕ is a graph isomorphism (there can be more than one) between the r_A -ball around x and the r_A -ball around $\phi(x)$, mapping x to $\phi(x)$.

Of course,

- $AB(x, y) = \sum_{z \in V} A(x, z)B(z, y)$
- $(A + B)(x, y) = A(x, y) + B(x, y)$
- $(\lambda A)(x, y) = \lambda(A(x, y))$
- $A^*(x, y) = \overline{A(y, x)}$

defines a $\star$ -algebra structure on P_G . Note that P_G is naturally represented on $L^2(V)$ by

$$A(f)(x) = \sum_{y \in V} A(x, y)f(y),$$

for $f \in L^2(V)$. These sort of operators were considered e.g. in [2], [3],[4].

In the case of tilings one can imbed P_G into a von Neumann algebra defined by the associated Delone system. In [2], we imbedded P_G into a continuous ring defined by the special construction of certain self-similar graphs. In this paper we define a von Neumann algebra N_G using the natural trace on P_G and consider the imbedding $\psi_G : P_G \rightarrow N_G$ (see Section 3). Now we can state the main results of this paper.

Theorem 1 *Let G be an AQ-graph, $B \in P_G$ a positive, self-adjoint operator (as an operator on $L^2(V)$). Suppose that for any $x, y \in V$, $B(x, y) \in \mathbb{Q}$. Then for any Følner-sequence $\{Q_n\}_{n=1}^\infty$, the spectral distribution functions N_{B_n} uniformly converge to the integrated density of states (IDS) function σ_B , where $B_n = p_{Q_n} B i_{Q_n}$ and σ_B is the von Neumann spectral function of the positive, self-adjoint element $\psi_G(B)$.*

Note that in [3], [4] similar uniform convergence results was proved for a class of AQ-graphs without the rationality assumption. In [2], we also considered certain AQ-graphs for which uniform convergence follows without the rationality assumption.

Theorem 2 *If G and B are as above then*

$$\int_0^\infty \log \lambda \, d\mu_B \geq 0,$$

where μ_B is the spectral measure associated to $\psi_G(B)$.

As a consequence of this theorem we have the following corollary:

Corollary 1.1 *If G and B are as above and λ is a point of discontinuity of the IDS function σ_B , then λ is an algebraic number.*

2 Uniform spectral convergence

In this section we prove some technicalities on the convergence of spectral distribution functions.

Proposition 2.1 *Let $\{f_n\}_{n=1}^\infty, f$ be monotone increasing right-continuous functions on the interval $[0, K]$ such that $f_n(K) = 1$ for any $n \geq 1$ and $f(K) = 1$ as well. Suppose that $\lim_{n \rightarrow \infty} f_n(\lambda) = f(\lambda)$ if λ is a point of continuity of f . Also suppose that if λ is a point of discontinuity of f , then*

$$\lim_{n \rightarrow \infty} J_n(\lambda) = J(\lambda),$$

where

$$J_n(\lambda) = f_n(\lambda) - f_n^-(\lambda) \quad J(\lambda) = f(\lambda) - f^-(\lambda)$$

are the jumps of our functions. Then $\{f_n\}_{n=1}^\infty$ uniformly converge to f .

Proof. First we need the following lemma.

Lemma 2.1

$$\lim_{n \rightarrow \infty} f_n^-(y) = f^-(y)$$

for any $y \in [0, K]$.

Proof. First suppose that

$$f^-(y) - \liminf_{n \rightarrow \infty} f_n^-(y) = \delta > 0. \tag{1}$$

Let $z < y$ be a continuity point of f such that $f^-(y) - f(z) < \frac{\delta}{2}$. Then for large enough n , $|f_n(z) - f(z)| < \frac{\delta}{2}$. Since $f_n(z) \leq f_n^-(y)$ (1) leads to a contradiction. Now suppose that

$$\limsup_{n \rightarrow \infty} f_n^-(y) - f^-(y) = \delta > 0. \quad (2)$$

Let $y < z$ a continuity point of f such that $f(z) - f(y) < \frac{\delta}{10}$. Again, for large enough n ;

$$|f_n(z) - f(z)| < \frac{\delta}{10} \quad \text{and} \quad |(f_n(y) - f_n^-(y)) - (f(y) - f^-(y))| < \frac{\delta}{10}.$$

Since $f_n(z) \geq f_n(y)$, (2) leads to a contradiction. ■

Now we turn to the proof of our proposition. If the uniform convergence does not exist, then there are points $\{x_{n_k}\}_{k=1}^\infty, y \in [0, K]$ such that $x_{n_k} \rightarrow y$ and

$$|f_{n_k}(x_{n_k}) - f(x_{n_k})| \geq \epsilon > 0 \quad (3)$$

Let $z < y$ be a continuity point of f such that $f^-(y) - f(z) < \frac{\epsilon}{10}$. If k is large enough, then

$$|f_{n_k}(z) - f(z)| < \frac{\epsilon}{10} \quad \text{and} \quad |f_{n_k}^-(y) - f^-(y)| < \frac{\epsilon}{10}.$$

Hence for large enough k :

$$|f_{n_k}(x_{n_k}) - f(x_{n_k})| < \epsilon,$$

if $z \leq x_{n_k} \leq y$. Now let $y < z$ be a continuity point of f such that $f(z) - f(y) < \frac{\epsilon}{10}$. Again if k is large, then

$$|f_{n_k}(z) - f(z)| < \frac{\epsilon}{10}, |f_{n_k}^-(y) - f^-(y)| < \frac{\epsilon}{10} \quad \text{and} \quad |f_{n_k}(y) - f(y)| < \frac{\epsilon}{10}.$$

Hence for large k , $|f_{n_k}(x_{n_k}) - f(x_{n_k})| < \epsilon$ if $y \leq x_{n_k} \leq z$. Hence (3) leads to a contradiction. ■

Suppose that μ_n, μ are probability measures on the interval $[0, K]$, and $\{\mu_n\}_{n=1}^\infty$ weakly converge to μ . Then if $F_n(\lambda) = \mu_n([0, \lambda]$, $F(\lambda) = \mu([0, \lambda]$, then $\lim_{n \rightarrow \infty} F_n(\lambda) = F(\lambda)$ if λ is a continuity point of F that is when $\mu(\{\lambda\}) = 0$. Suppose that

- μ_n weakly converge to μ and
- if $\mu(\{\lambda\}) > 0$, then $\mu_n(\{\lambda\}) \rightarrow \mu(\{\lambda\})$.

Then by our Proposition, $\{F_n\}_{n=1}^\infty$ converge to F uniformly.

Remark: Consider the situation in Statement 1. If $B \in \mathbb{C}\Gamma$ is a positive, self-adjoint element, then by [1]

$$\mathrm{Tr}_\Gamma S_\lambda^B = \lim_{n \rightarrow \infty} \frac{\dim_{\mathbb{C}} \mathrm{Ker}(B - \lambda)}{|F_n|},$$

where S_λ^B is the orthogonal projection onto the λ -eigenspace of B . Hence by Proposition 2.1, N_{B_n} uniformly converge to the spectral measure σ_B (the pointwise convergence was considered in [7]).

3 The pattern-frequency algebra construction

Let G be an AQ-graph and P_G be the $\star$ -algebra of pattern-invariant matrices. Then for $A \in P_G$ we define

$$\mathrm{Tr}_G(A) := \lim \frac{1}{|Q_n|} \sum_{x \in Q_n} A(x, x), \quad (4)$$

where $\{Q_n\}_{n=1}^\infty$ is a Følner-sequence in G .

Proposition 3.1 (a) *The limit in (4) exists.*

(b) *$\mathrm{Tr}_G(A)$ does not depend on the choice of the Følner-sequence.*

(c) *Tr_G is a trace, that is a linear functional satisfying $\mathrm{Tr}_G(AB) = \mathrm{Tr}_G(BA)$.*

(d) *Tr_G is faithful, that is $\mathrm{Tr}_G(A^*A) > 0$ if $A \neq 0$.*

Proof. Note that

$$\frac{1}{|Q_n|} \sum_{x \in Q_n} A(x, x) = \frac{1}{|Q_n|} \sum_{\alpha \in P_{r_A}(G)} |Q_n^\alpha| A_\alpha,$$

where Q_n^α is the number of vertices in Q_n with r_A -pattern α and $A_\alpha = A(x, x)$ if x has r_A -pattern α . Since $\lim_{n \rightarrow \infty} \frac{|Q_n^\alpha|}{|Q_n|} = P(\alpha)$, (a) and (b) immediately follows.

Now let $A, B \in P_G$, then

$$\mathrm{Tr}_G(AB) = \lim_{n \rightarrow \infty} \left(\frac{1}{|Q_n|} \sum_{x \in Q_n} \sum_{y \in V} A(x, y) B(y, x) \right)$$

$$\mathrm{Tr}_G(BA) = \lim_{n \rightarrow \infty} \left(\frac{1}{|Q_n|} \sum_{x \in Q_n} \sum_{y \in V} B(x, y) A(y, x) \right)$$

That is

$$|\mathrm{Tr}_G(AB) - \mathrm{Tr}_G(BA)| \leq \lim_{n \rightarrow \infty} \frac{2}{|Q_n|} \sum_{x \in \partial_D Q_n} m_A m_B, \quad (5)$$

where $m_A = \sup_{x, y} |A(x, y)|$, $m_B = \sup_{x, y} |B(x, y)|$, and

$$\partial_D Q_n := \{x \in Q_n \mid \text{there exists } y \notin Q_n, d_G(x, y) \leq \max(r_A, r_B)\}.$$

Note that $m_A, m_B < \infty$ by the pattern-invariance. Also, $\lim_{n \rightarrow \infty} \frac{|\partial_D Q_n|}{|Q_n|} = 0$ by the Følner-property and the bounded vertex degree condition. Hence by (5) (c) follows. Finally, let $0 \neq A \in P_G$. Then

$$\mathrm{Tr}_G(A^*A) = \frac{1}{|Q_n|} \sum_{x \in Q_n} \left(\sum_{y \in V} A(x, y) \overline{A(y, x)} \right) = \frac{1}{|Q_n|} \sum_{x \in Q_n} \left(\sum_{y \in V} |A(x, y)|^2 \right).$$

That is $\mathrm{Tr}_G(A^*A) = \sum_{\alpha \in P_{r_A}(G)} P(\alpha) L_\alpha$, where $L_\alpha = \sum_{y \in V} |A(x, y)|^2$ if x has the r_A -pattern α . Therefore (d) follows. $\blacksquare$

Now the pattern-frequency algebra is constructed by the GNS-construction the usual way. The algebra P_G is a pre-Hilbert space with respect to the inner product $\langle A, B \rangle = \mathrm{Tr}_G(B^*A)$. Then $L_A B = AB$ defines a representation of P_G on this pre-Hilbert space.

Lemma 3.1 *L_A is a bounded operator if $A \in P_G$.*

Proof. Let $A, B \in P_G$. Denote by $\| \cdot \|$ the pre-Hilbert space norm. Then

$$\|B\|^2 = \lim_{n \rightarrow \infty} \left(\frac{1}{|Q_n|} \sum_{x \in Q_n} \left(\sum_{y \in V} |B(x, y)|^2 \right) \right).$$

$$\|L_A B\|^2 = \lim_{n \rightarrow \infty} \left(\frac{1}{|Q_n|} \sum_{x \in Q_n} \left(\sum_{y \in V} BB^*(x, y) A^* A(y, x) \right) \right) \leq \lim_{n \rightarrow \infty} \left(\frac{K}{|Q_n|} \sum_{x \in Q_n} \left(\sum_{y \in V} |BB^*(x, y)| \right) \right),$$

where $K = \sup_{x, y} |A^* A(x, y)|$. Note that

$$\begin{aligned} |BB^*(x, y)| &= \left| \sum_{z \in V} B(x, z) B^*(z, y) \right| \leq \sum_{z \in V} |B(x, z)| |B(y, z)| \\ &\leq \frac{1}{2} \sum_{z \in V} (|B(x, z)|^2 + |B(y, z)|^2). \end{aligned}$$

Let $t_x := \sum_{y \in V} |B(x, y)|^2$. Then

$$\|B\|^2 = \lim_{n \rightarrow \infty} \frac{1}{|Q_n|} \sum_{x \in Q_n} t_x.$$

$$\|L_A B\|^2 = \lim_{n \rightarrow \infty} \left(\frac{1}{|Q_n|} \sum_{x \in Q_n} \left(\sum_{y \in V} \frac{1}{2} (t_x + t_y) \right) \leq \lim_{n \rightarrow \infty} \frac{KM}{|Q_n|} \sum_{x \in Q_n} t_x, \right.$$

where M is the maximal number of vertices in an r_{A^*A} -ball of G . Hence $\|L_A B\|^2 \leq KM\|B\|^2$. ■

Thus P_G is represented by bounded operators on the Hilbert-space closure. Now the pattern-frequency algebra N_G is defined as the weak-closure of P_G . then $\text{Tr}_G(A) = \langle L_A 1, 1 \rangle$ extends to N_G as an ultraweakly continuous, faithful trace. The finite von Neumann algebra N_G shall play the role of the group von Neumann algebra in our paper.

4 Spectral approximation

Proposition 4.1 *If B is a positive, self-adjoint operator in P_G , then $\psi_G(B)$ is positive self-adjoint as an element of N_G .*

Proof. The self-adjointness is obvious, since the representation $\psi_G : P_G \rightarrow N_G$ preserves the $*$ -structure:

$$\langle L_B X, Y \rangle = \text{Tr}_G(Y^* B X)$$

$$\langle X, L_B Y \rangle = \text{Tr}_G((BY)^* X) = \text{Tr}_G(Y^* B X).$$

The operator B is positive in P_G if and only if $\langle B(f), f \rangle_{L^2(V)} \geq 0$ for any $f \in L^2(V)$. That is

$$\sum_{y, z \in V} B(y, z) f(z) \overline{f(y)} \geq 0. \quad (6)$$

On the other hand, $\psi_G(B)$ is positive if $\text{Tr}_G(U^* B U) \geq 0$ for any $U \in P_G$.

$$\begin{aligned} \text{Tr}_G(U^* B U) &= \lim_{n \rightarrow \infty} \left(\frac{1}{|Q_n|} \sum_{x \in Q_n} \sum_{y, z \in V} U^*(x, y) B(y, z) U(z, x) \right) = \\ &= \lim_{n \rightarrow \infty} \left(\frac{1}{|Q_n|} \sum_{x \in Q_n} \sum_{y, z \in V} B(y, z) U(z, y) \overline{U(y, x)} \right). \end{aligned}$$

By (6), for any $x \in Q_n$:

$$\sum_{y,z \in V} B(y,z)U(z,y)\overline{U(y,x)} \geq 0.$$

Hence the positivity follows. $\blacksquare$

Now we need some estimates for the norms of the finite dimensional linear transforms B_n .

Proposition 4.2 *If $B \in P_G$ is positive, self-adjoint operator then B_n is positive, self-adjoint as well. Furthermore, there exists $K > 0$ such that $\|B_n\| \leq K$, for any $n \geq 1$, where the norm of B_n is considered as a linear transformation in $\text{Mat}_{Q_n \times Q_n}(\mathbb{C})$.*

Proof. Recall that $B_n = p_{Q_n} B p_{Q_n}^*$, hence the positivity and self-adjointness are obvious. The norm-estimate immediately follows from the next lemma.

Lemma 4.1 *Let $H(V, E)$ be a finite graph, where the vertex degrees are not greater than d . Suppose that $A : V \times V \rightarrow \mathbb{C}$ is a matrix such that*

- $|A(x, y)| \leq m$
- $A(x, y) = 0$ if $d_H(x, y) > r$.

Then $\|A\| \leq C_{d,m,r}$, where the constant $C_{d,m,r} > 0$ depends only on the parameters, not on the graph H itself.

Proof. For $f, g \in L^2(V)$ unit vectors,

$$\begin{aligned} |\langle A(f), g \rangle| &= \left| \sum_{x,y \in V} A(x,y) f(x) \overline{g(y)} \right| \leq m \left| \sum_{x,y \in V, d_H(x,y) \leq r} f(x) \overline{g(y)} \right| \leq \\ &\leq m \sum_{x,y \in V, d_H(x,y) \leq r} (|f(x)|^2 + |g(y)|^2). \end{aligned}$$

The number of occurrences of $|g(y)|^2$ on the right hand side is not greater than the maximal possible number $T(r, d)$ of vertices in a ball of radius r in a graph of vertex degrees bounded by d . This quantity depends only on r and d . Thus

$$\|A\| = \sup_{f,g \in L^2(V), \|f\|=1, \|g\|=1} |\langle A(f), g \rangle| \leq m(1 + T(r, d)). \quad \blacksquare$$

Now we prove our key trace approximation formula.

Proposition 4.3 *Let $B \in P_G$, then for any $k \geq 1$,*

$$\text{Tr}_G(B^k) = \lim_{n \rightarrow \infty} \frac{1}{|Q_n|} \text{Tr}(B_n^k).$$

Proof.

$$\begin{aligned} \text{Tr}_G(B^k) &= \lim_{n \rightarrow \infty} \frac{1}{|Q_n|} \sum_{x \in Q_n} B^k(x, x) = \\ &= \lim_{n \rightarrow \infty} \frac{1}{|Q_n|} \sum_{x \in Q_n} \sum_{y_1, y_2, \dots, y_{k-1} \in V} B(x, y_1) B(y_1, y_2) \dots B(y_{k-1}, x). \end{aligned}$$

On the other hand,

$$\frac{1}{|Q_n|} \text{Tr}(B_n^k) = \frac{1}{|Q_n|} \sum_{x \in Q_n} \sum_{y_1, y_2, \dots, y_{k-1} \in Q_n} B(x, y_1) B(y_1, y_2) \dots B(y_{k-1}, x).$$

Note that if $x, y \in Q_n$ then $B(x, y) = B_n(x, y)$. Clearly,

$$\begin{aligned} & \left| \frac{1}{|Q_n|} \sum_{x \in Q_n} \sum_{y_1, y_2, \dots, y_{k-1} \in V} B(x, y_1) B(y_1, y_2) \dots B(y_{k-1}, x) - \right. \\ & \left. - \frac{1}{|Q_n|} \sum_{x \in Q_n} \sum_{y_1, y_2, \dots, y_{k-1} \in Q_n} B(x, y_1) B(y_1, y_2) \dots B(y_{k-1}, x) \right| \leq \frac{1}{|Q_n|} m_B^k S_n, \end{aligned}$$

where $m_B = \sup_{x, y \in V} B(x, y)$ and S_n is the set of strings $\{x, y_1, y_2, \dots, y_{k-1}\}$ such that

- $d_G(x, y_1) \leq r$.
- $d_G(y_i, y_{i+1}) \leq r$ for any $1 \leq i \leq k-2$.
- For some $1 \leq i \leq k-1$: $y_i \notin Q_n$.

Therefore our proposition follows from the lemma below.

Lemma 4.2 $\lim_{n \rightarrow \infty} \frac{|S_n|}{|Q_n|} = 0$.

Proof. For $a \in \mathbb{N}$, let again

$$\partial_a Q_n := \{x \in Q_n \mid \text{there exists } y \notin Q_n, d_G(x, y) \leq a\}.$$

By the Følner-property and the vertex degree condition $\lim_{n \rightarrow \infty} \frac{|\partial_a Q_n|}{|Q_n|} = 0$. If x is a starting vertex of a string in S_n , then $x \in \partial_{kr} Q_n$. The number of choices for y_1 is not greater than q_r , where q_r is the maximal number of vertices in an r -ball of G . Hence

$$|S_n| \leq |\partial_{kr} Q_n| q_r^{k-1}.$$

Thus our lemma follows. ■

5 The integrated density of states

Let G be an AQ-graph, $B \in P_G$ be positive, self-adjoint and $\{B_n\}_{n=1}^\infty$ be the associated, finite dimensional linear transformations. Since $\psi_G(B) \in N_G$ is positive, self-adjoint element of a von Neumann-algebra we have the spectral theorem:

$$\psi_G(B) := \int_0^\infty \lambda dE_\lambda^B,$$

where E_λ^B are the associated spectral projections $\chi_{[0,\lambda]}(\psi_G(B))$. The spectral measure is defined by

$$\sigma_B(\lambda) = \mu_B[0, \lambda] := \text{Tr}_G(E_\lambda^B).$$

For the finite dimensional operators B_n

$$N_{B_n}(\lambda) = \frac{1}{|Q_n|} \text{Tr}(E_\lambda^{B_n})$$

are the usual spectral distributions. Let $K > 0$ be a real number such that $\|B\|, \|B_n\| < K$.

Then if $P \in \mathbb{R}[x]$ is a polynomial,

$$\int_0^K P(\lambda) d\mu_{B_n}(\lambda) = \frac{1}{|Q_n|} \text{Tr}(P(B_n)).$$

$$\int_0^K P(\lambda) d\mu_B(\lambda) = \text{Tr}_G(P(B_n)).$$

Hence by Proposition 4.3

$$\lim_{n \rightarrow \infty} \int_0^K P(\lambda) d\mu_{B_n}(\lambda) = \int_0^K P(\lambda) d\mu_B(\lambda).$$

That is the probability measures μ_{B_n} weakly converge to μ_B . In other words, σ_B is the integrated density of states. This immediately implies the associated Shubin-formula:

$$\lim_{n \rightarrow \infty} N_{B_n}(\lambda) = \text{Tr}_G E_\lambda^B,$$

for any $\lambda \geq 0$ continuity point of σ_B .

6 Approximation of the ground state density

The goal of this section is to prove the following proposition.

Proposition 6.1 *If B is a positive, self-adjoint element of P_G and $B(x, y)$ is a rational number for any $x, y \in V$, then*

$$\lim_{n \rightarrow \infty} \frac{\dim_{\mathbb{C}} \text{Ker } B_n}{|Q_n|} = \text{Tr}_G E_0^B.$$

Proof. The proof is very similar to Theorem 6.9 [7] nevertheless there are some details we would like to make precise for the reader who is not familiar with original paper of Lück [5]. Note that B has only finitely many different values, therefore we can suppose (after multiplying by an appropriate integer) that B has only integer values.

First we need some notations. For a monotone function f ,

$$f^+(\lambda) = \inf_{\epsilon \rightarrow 0} f(\lambda + \epsilon)$$

$$\overline{\sigma_B}(\lambda) := \limsup_{n \rightarrow \infty} N_{B_n}(\lambda)$$

$$\underline{\sigma_B}(\lambda) := \liminf_{n \rightarrow \infty} N_{B_n}(\lambda).$$

Let us recall Lemma 6.5 [7] in the special case we need it.

Lemma 6.1 *For any polynomial $P_k \in \mathbb{R}[x]$ which has the property $\chi_{[0, \lambda]}(x) \leq P_k(x) \leq \frac{1}{k} \chi_{[0, \lambda]}(x) + \chi_{[0, \lambda + \frac{1}{k}]}(x)$ for any $0 \leq x \leq K$, we have the following inequalities:*

$$\sigma_B(\lambda) \leq \text{Tr}_G(P_k(\psi_G(B))) \leq \sigma_B(\lambda + \frac{1}{k}) + \frac{1}{k}.$$

$$N_{B_n}(\lambda) \leq \frac{1}{|Q_n|} \text{Tr}(P_k(B_n)) \leq N_{B_n}(\lambda + \frac{1}{k}) + \frac{1}{k}.$$

We also need the appropriate version of Proposition 6.7 [7].

Proposition 6.2

$$\overline{\sigma_B}(\lambda) \leq \sigma_B(\lambda) = \underline{\sigma_B}^+(\lambda) = \overline{\sigma_B}^+(\lambda).$$

Proof. First choose the polynomial P_k as in the previous lemma. Then

$$N_{B_n}(\lambda) \leq \frac{1}{|Q_n|} \text{Tr}(P_k(B_n)) \leq N_{B_n}(\lambda + \frac{1}{k}) + \frac{1}{k}.$$

That is as $n \rightarrow \infty$:

$$\overline{\sigma_B}(\lambda) \leq \text{Tr}_G P_k(\psi_G(B)) \leq \underline{\sigma_B}(\lambda + \frac{1}{k}) + \frac{1}{k}.$$

Now we let $k \rightarrow \infty$. Then:

$$\lim_{k \rightarrow \infty} \text{Tr}_G P_k(\psi_G(B)) = \lim_{k \rightarrow \infty} \langle P_k(\psi_G(B))(1), 1 \rangle = \langle \chi_{[0, \lambda]}(\psi_G(B))(1), 1 \rangle = \text{Tr}_G E_\lambda^B,$$

since $\{P_k(\psi_G(B))\}_{k=1}^\infty$ strongly converge to $\chi_{[0, \lambda]}(\psi_G(B))$. Therefore,

$$\overline{\sigma_B}(\lambda) \leq \sigma_B(\lambda) \leq \underline{\sigma_B}^+(\lambda).$$

Clearly,

$$\sigma_B(\lambda) \leq \underline{\sigma_B}(\lambda + \epsilon) \leq \overline{\sigma_B}(\lambda + \epsilon) \leq \sigma_B(\lambda + \epsilon).$$

That is

$$\sigma_B(\lambda) = \underline{\sigma_B}^+(\lambda) = \overline{\sigma_B}^+(\lambda).$$

Now we need a lemma on Riemann-Stieltjes integral. The proof can be found in [6] in Lemma 4.1.

Lemma 6.2 *Let f be a continuously differentiable function on the positive reals and μ be a probability measure on the $[0, K]$ interval. Suppose that F is the distribution function of μ , that is $\mu[0, \lambda] = F(\lambda)$. Then for any $0 < \epsilon \leq K$:*

$$\int_\epsilon^K f(\lambda) d\mu = - \int_\epsilon^K f'(\lambda) F(\lambda) d\lambda + f(K)F(K) - f(\epsilon)F(\epsilon).$$

Let $|det|_1(B_n)$ is defined as the product of non-negative eigenvalues of B_n , this is the lowest nonzero coefficient in the characteristic polynomial of B_n hence a non-zero integer.

Therefore,

$$\log \det_{Q_n}(B_n) := \frac{1}{|Q_n|} \log(|det|_1(B_n)) \geq 0. \quad (7)$$

By Lemma 6.2,

$$\log \det_{Q_n}(B_n) = \log K - \log \epsilon \cdot N_{B_n}(\epsilon) - \int_\epsilon^K N_{B_n}(\lambda) \frac{1}{\lambda} d\lambda,$$

where $\epsilon > 0$ is a real number below the smallest eigenvalue of B_n .

Since $\int_\epsilon^K \frac{N_{B_n}(\lambda)}{\lambda} d\lambda = (\log K - \log \epsilon) N_{B_n}(0)$:

$$\log \det_{Q_n}(B_n) = \log K(1 - N_{B_n}(0)) - \int_0^K \frac{N_{B_n}(\lambda) - N_{B_n}(0)}{\lambda} d\lambda. \quad (8)$$

Hence by (7)

$$\int_0^K \frac{N_{B_n}(\lambda) - N_{B_n}(0)}{\lambda} d\lambda \leq \log K. \quad (9)$$

Now we estimate $\int_0^K \frac{\sigma_B(\lambda) - \sigma_B(0)}{\lambda} d\lambda$.

$$\int_\epsilon^K \frac{\sigma_B(\lambda) - \sigma_B(0)}{\lambda} d\lambda \leq \int_\epsilon^K \frac{\underline{\sigma}_B^+(\lambda) - \sigma_B(0)}{\lambda} d\lambda \leq \int_\epsilon^K \frac{\underline{\sigma}_B(\lambda) - \sigma_B(0)}{\lambda} d\lambda.$$

Furthermore,

$$\begin{aligned} \int_\epsilon^K \frac{\underline{\sigma}_B(\lambda) - \sigma_B(0)}{\lambda} d\lambda &\leq \int_\epsilon^K \frac{\underline{\sigma}_B(\lambda) - \overline{\sigma}_B(0)}{\lambda} d\lambda = \int_\epsilon^K \frac{\liminf_{n \rightarrow \infty} N_{B_n}(\lambda) - N_{B_n}(0)}{\lambda} d\lambda \leq \\ &\leq \liminf_{n \rightarrow \infty} \int_\epsilon^K \frac{N_{B_n}(\lambda) - N_{B_n}(0)}{\lambda} d\lambda \leq \log K. \end{aligned}$$

Now we take the limit $\epsilon \rightarrow 0$:

$$\begin{aligned} &\int_0^K \frac{\sigma_B(\lambda) - \sigma_B(0)}{\lambda} d\lambda \leq \\ &\leq \int_0^K \frac{\underline{\sigma}_B(\lambda) - \overline{\sigma}_B(0)}{\lambda} d\lambda \leq \sup_{\epsilon \rightarrow 0} \liminf_{n \rightarrow \infty} \frac{N_{B_n}(\lambda) - N_{B_n}(0)}{\lambda} d\lambda \leq \log K. \end{aligned} \quad (10)$$

Therefore $\lim_{\lambda \rightarrow 0} \underline{\sigma}_B(\lambda) = \overline{\sigma}_B(0)$. That is by Proposition 6.2: $\overline{\sigma}_B(0) = \sigma_B(0)$, hence

$$\limsup_{n \rightarrow \infty} N_{B_n}(0) = \sigma_B(0). \quad (11)$$

Since one can consider any subsequence of B_n , (11) implies that

$$\lim_{n \rightarrow \infty} N_{B_n}(0) = \lim_{n \rightarrow \infty} \frac{\dim_{\mathbb{C}} \text{Ker } B_n}{|Q_n|} = \sigma_B(0) = \text{Tr}_G E_0^B. \quad \blacksquare$$

7 The proof of Theorem 2

We would like to show that $\int_0^\infty \log \lambda d\mu_B \geq 0$. Note that

$$\int_0^\infty \log \lambda d\mu_B = \lim_{\epsilon \rightarrow 0} \int_\epsilon^K \log \lambda d\mu_B = \lim_{\epsilon \rightarrow 0} (\log K(1 - \sigma_B(\epsilon)) - \int_\epsilon^K \frac{\sigma_B(\lambda) - \sigma_B(0)}{\lambda} d\lambda).$$

Hence

$$\int_0^\infty \log \lambda d\mu_B = \log K(1 - \sigma_B(0)) - \int_0^K \frac{\sigma_B(\lambda) - \sigma_B(0)}{\lambda} d\lambda.$$

By (10)

$$\begin{aligned} \sup_{\epsilon > 0} \liminf_{n \rightarrow \infty} \int_{\epsilon}^K \frac{N_{B_n}(\lambda) - N_{B_n}(0)}{\lambda} d\lambda &\leq \liminf_{n \rightarrow \infty} \sup_{\epsilon > 0} \int_{\epsilon}^K \frac{N_{B_n}(\lambda) - N_{B_n}(0)}{\lambda} d\lambda = \\ &= \liminf_{n \rightarrow \infty} \int_0^K \frac{N_{B_n}(\lambda) - N_{B_n}(0)}{\lambda} d\lambda \leq \log K(1 - N_{B_n}(0)). \end{aligned}$$

Thus,

$$\begin{aligned} \log \det(B) &= \log K(1 - \sigma_B(0)) - \int_0^K \frac{\sigma_B(\lambda) - \sigma_B(0)}{\lambda} d\lambda \geq \\ &= \log k(1 - \lim_{n \rightarrow \infty} N_{B_n}(0)) - \liminf_{n \rightarrow \infty} \int_0^K \frac{N_{B_n}(\lambda) - N_{B_n}(0)}{\lambda} d\lambda = \\ &= \limsup_{n \rightarrow \infty} (\log(K)(1 - \lim_{n \rightarrow \infty} N_{B_n}(0)) - \int_0^K \frac{N_{B_n}(\lambda) - N_{B_n}(0)}{\lambda} d\lambda) = \\ &= \limsup_{n \rightarrow \infty} \log \det_{Q_n}(B_n) \geq 0. \quad \blacksquare \end{aligned}$$

8 The proof of Theorem 1

By Proposition 2.1, it is enough to prove that

$$\lim_{n \rightarrow \infty} \frac{\dim_{\mathbb{C}} \text{Ker}(B_n - \lambda)}{|Q_n|} = \text{Tr}_G S_{\lambda}^B,$$

where S_{λ}^B is the orthogonal projection onto the λ -eigenspace of $\psi_G(B)$. Observe that $S_{\lambda}^B = S_0^{(B-\lambda)^2}$. Therefore by Proposition 6.1, Theorem 1 follows from the lemma below.

Lemma 8.1

$$\lim_{n \rightarrow \infty} \frac{\dim_{\mathbb{C}} \text{Ker}(B_n - \lambda) - \dim_{\mathbb{C}} \text{Ker}(p_{Q_n}(B_{\lambda})^2 i_{Q_n})}{|Q_n|} = 0.$$

Proof. Let $Q'_n = Q_n \setminus \partial_{2r_{B_n}} Q_n$. Observe that if $\text{supp} f \subseteq Q'_n$, then

$$(p_{Q_n}(B_{\lambda})^2 i_{Q_n})f = (p_{Q_n}(B_{\lambda}) i_{Q_n})^2 f.$$

Hence

$$\text{Ker}(B_n - \lambda) \cap L^2(Q'_n) = \text{Ker}(p_{Q_n}(B_{\lambda})^2 i_{Q_n}) \cap L^2(Q'_n).$$

Since $\lim_{n \rightarrow \infty} \frac{|Q'_n|}{|Q_n|} = 1$ the lemma follows. $\blacksquare$

References

- [1] J. DODZIUK, P. LINNELL, V. MATHAI, T. SCHICK AND S. YATES, Approximating L^2 -invariants and the Atiyah conjecture, *Comm. Pure Appl. Math.* **56** (2003) no.7 839-873.
- [2] G. ELEK, Aperiodic order, integrated density of states and the continuous algebras of John von Neumann, *preprint* URL: <http://www.arxiv.org/math-ph/0606061>
- [3] D. LENZ AND P. STOLLMANN, Aperiodic order and quasicrystals: spectral properties. *Ann. Henri Poincaré*, **4** (2003), suppl2. 933-942.
- [4] D. LENZ AND P. STOLLMANN, Ergodic theorem for Delone dynamical systems and existence of the integrated density of states. *to appear in Journal d' Anal. Math.*
- [5] W. LÜCK, Approximating L^2 -invariants by their finite-dimensional analogues. *Geom. Funct. Anal.* **4** (1994), no. 4, 455-481.
- [6] W. LÜCK, L^2 -torsion and 3-manifolds. *Low-dimensional topology (Knoxville, TN, 1992)*, 75-107. Conf. Proc. Lecture Notes Geom. Topology, III, Internat. Press, Cambridge, MA, 1994.
- [7] T. SCHICK, L^2 -determinant class and approximation of L^2 -Betti numbers *Trans. Amer. Math. Soc.* **353** (2001) no. 8, 3247-3265.